

A class of digit extraction BBP-type formulas in general binary bases*

Kunle Adegoke^{†1}, Jaume Oliver Lafont² and Olawanle Layeni³

¹Department of Physics and Engineering Physics,
Obafemi Awolowo University, Ile-Ife, 220005 Nigeria

²Conselleria d'Educacio i Cultura, Govern de les Illes Balears,
Palma, Spain

³Department of Mathematics,
Obafemi Awolowo University, Ile-Ife, 220005 Nigeria

Abstract

BBP-type formulas are usually discovered experimentally, one at a time and in specific bases, through computer searches. In this paper, however, we derive directly, without doing any searches, explicit digit extraction BBP-type formulas in general binary bases $b = 2^{12p}$, for p positive odd integers. As particular examples, new binary formulas are presented for $\pi\sqrt{3}$, $\pi\sqrt{3}\log 2$, $\sqrt{3}\operatorname{Cl}_2(\pi/3)$ and a couple of other polylogarithm constants. A variant of the formula for $\pi\sqrt{3}\log 2$ derived in this paper has been known for over ten years but was hitherto unproved. Binary BBP-type formulas for the logarithms of an infinite set of primes and binary BBP-type representations for the arctangents of an infinite set of rational numbers are also presented. Finally, new binary BBP-type zero relations are established.

*MSC 2010: 11Y60, 30B99

[†]adegoke00@gmail.com

Contents

1	Introduction	2
2	Definitions and Notation	3
3	Degree s Formulas	4
4	Degree 1 Formulas	7
5	Degree 2 Formulas	10
6	Interesting Particular Cases	12
6.1	Degree 1 Binary BBP-type Formulas	12
6.1.1	Binary BBP-type formula for $\log 2$	13
6.1.2	Binary BBP-type formula for $\pi\sqrt{3}$	13
6.1.3	Binary BBP-type formula for $\sqrt{3}\log(2 + \sqrt{3})$	13
6.1.4	Binary BBP-type formula for $\arctan 1/6$	14
6.2	Degree 2 Binary BBP-type Formulas	14
6.2.1	Binary BBP-type formula for $\sqrt{3}\text{Cl}_2(\pi/3)$	15
6.2.2	Binary BBP-type formula for $\pi\sqrt{3}\log 2$	15
6.2.3	Binary BBP-type formula for Catalan's constant G	16
6.3	Binary Zero Relations	16
7	Conclusion	18

1 Introduction

BBP-type formulas are formulas of the form

$$\alpha = \sum_{k=0}^{\infty} 1/b^k \sum_{j=1}^l a_j / (kl + j)^s$$

where s , b , l (degree, base, length respectively) and a_j are integers, and α is some constant. Formulas of this type were first introduced in a 1996 paper [1], where a formula of this type for π was given. Such formulas allow digit extraction — the i -th digit of a mathematical constant α in base b can be calculated directly, without needing to compute any of the previous $i - 1$

digits, by means of simple algorithms that do not require multiple-precision arithmetic [2].

Apart from digit extraction, another reason the study of BBP-type formulas has continued to attract attention is that BBP-type constants are conjectured to be either rational or normal to base b [3, 4, 5], that is their base- b digits are randomly distributed.

BBP-type formulas are usually discovered experimentally, one at a time and in specific bases, through computer searches. In this paper, however, we derive explicit digit extraction BBP-type formulas in general binary bases $b = 2^{12p}$, for $p \in \mathbb{Z}^+$ and $\text{mod}(p, 2) = 1$.

2 Definitions and Notation

The polylogarithm functions denoted by Li in this paper are defined by

$$\text{Li}_s[z] = \sum_{k=1}^{\infty} \frac{z^k}{k^s}, \quad |z| \leq 1, s \in \mathbb{Z}^+.$$

For $|z| = 1$ and $x \in [0, 2\pi]$ we have

$$\begin{aligned} \text{Li}_{2n}[e^{ix}] &= \text{Gl}_{2n}(x) + i\text{Cl}_{2n}(x) \\ \text{Li}_{2n+1}[e^{ix}] &= \text{Cl}_{2n+1}(x) + i\text{Gl}_{2n+1}(x), \end{aligned}$$

where Gl and Cl are Clausen sums [6] defined, for $n \in \mathbb{Z}^+$ by

$$\begin{aligned} \text{Cl}_{2n}(x) &= \sum_{k=1}^{\infty} \frac{\sin kx}{k^{2n}}, & \text{Cl}_{2n+1}(x) &= \sum_{k=1}^{\infty} \frac{\cos kx}{k^{2n+1}} \\ \text{Gl}_{2n}(x) &= \sum_{k=1}^{\infty} \frac{\cos kx}{k^{2n}}, & \text{Gl}_{2n+1}(x) &= \sum_{k=1}^{\infty} \frac{\sin kx}{k^{2n+1}}. \end{aligned}$$

We shall find the following formulas useful:

$$\begin{aligned} \text{Gl}_{2n}(x) &= (-1)^{1+[n/2]} 2^{n-1} \pi^n \text{B}_n(x/2\pi)/n! \\ \frac{1}{m^{n-1}} \text{Cl}_n(mx) &= \sum_{r=0}^{m-1} \text{Cl}_n(x + 2\pi r/m). \end{aligned}$$

Here $[n/2]$ denotes the integer part of $n/2$ and B_n are the Bernoulli polynomials defined by

$$\frac{te^{xt}}{e^t - 1} = \sum_{n=0}^{\infty} \frac{B_n(x)t^n}{n!}.$$

In order to save space, we will give the BBP-type formulas using the compact P-notation [2]:

$$P(s, b, l, A) \equiv \sum_{k=0}^{\infty} \frac{1}{b^k} \sum_{j=1}^l \frac{a_j}{(kl + j)^s}, \quad (2.1)$$

where s , b and l are integers, and $A = (a_1, a_2, \dots, a_l)$ is a vector of integers.

3 Degree s Formulas

Using the identities

$$\operatorname{Re} \operatorname{Li}_s \left[\frac{1}{\sqrt{2}^p} \exp ix \right] = \sum_{k=1}^{\infty} \left[\frac{1}{\sqrt{2}^{pk}} \frac{1}{k^s} \cos kx \right] \quad (3.1)$$

and

$$\operatorname{Im} \operatorname{Li}_s \left[\frac{1}{\sqrt{2}^p} \exp ix \right] = \sum_{k=1}^{\infty} \left[\frac{1}{\sqrt{2}^{pk}} \frac{1}{k^s} \sin kx \right] \quad (3.2)$$

for $x \in \{\pi/12, 5\pi/12, 7\pi/12, 11\pi/12\}$ and the fact that

$$\cos \left(\frac{\pi}{12} \right) = \frac{\sqrt{3} + 1}{2\sqrt{2}} = \sin \left(\frac{7\pi}{12} \right) = \sin \left(\frac{5\pi}{12} \right) = -\cos \left(\frac{11\pi}{12} \right)$$

and

$$\sin \left(\frac{\pi}{12} \right) = \frac{\sqrt{3} - 1}{2\sqrt{2}} = \cos \left(\frac{5\pi}{12} \right) = \sin \left(\frac{11\pi}{12} \right) = -\cos \left(\frac{7\pi}{12} \right),$$

it is not difficult to obtain the following results, written in the P-notation ((2.1)):

$$\begin{aligned}
& \operatorname{Re} \operatorname{Li}_s \left[\frac{1}{\sqrt{2}^p} \exp \left(\frac{i\pi}{12} \right) \right] + \operatorname{Re} \operatorname{Li}_s \left[\frac{1}{\sqrt{2}^p} \exp \left(\frac{7i\pi}{12} \right) \right] \\
&= \frac{1}{2^{12p}} P(s, 2^{12p}, 24, (2^{(-\frac{1}{2}+\frac{p}{2}+11p)}, 0, 2^{(\frac{1}{2}-\frac{p}{2}+11p)}, 2^{10p}, \\
&\quad -2^{(-\frac{1}{2}+\frac{p}{2}+9p)}, 0, 2^{(-\frac{1}{2}+\frac{p}{2}+8p)}, -2^{8p}, -2^{(\frac{1}{2}-\frac{p}{2}+8p)}, 0, -2^{(-\frac{1}{2}+\frac{p}{2}+6p)}, \\
&\quad -2^{1+6p}, -2^{(-\frac{1}{2}+\frac{p}{2}+5p)}, 0, -2^{(\frac{1}{2}-\frac{p}{2}+5p)}, -2^{4p}, 2^{(-\frac{1}{2}+\frac{p}{2}+3p)}, 0, \\
&\quad -2^{(-\frac{1}{2}+\frac{p}{2}+2p)}, 2^{2p}, 2^{(\frac{1}{2}-\frac{p}{2}+2p)}, 0, 2^{(-\frac{1}{2}+\frac{p}{2})}, 2)) , \tag{3.3}
\end{aligned}$$

$$\begin{aligned}
& \operatorname{Re} \operatorname{Li}_s \left[\frac{1}{\sqrt{2}^p} \exp \left(\frac{i\pi}{12} \right) \right] - \operatorname{Re} \operatorname{Li}_s \left[\frac{1}{\sqrt{2}^p} \exp \left(\frac{7i\pi}{12} \right) \right] \\
&= \frac{\sqrt{3}}{2^{12p}} P(s, 2^{12p}, 24, (2^{(-\frac{1}{2}+\frac{p}{2}+11p)}, 2^{11p}, 0, 0, \\
&\quad 2^{(-\frac{1}{2}+\frac{p}{2}+9p)}, 0, -2^{(-\frac{1}{2}+\frac{p}{2}+8p)}, 0, 0, -2^{7p}, -2^{(-\frac{1}{2}+\frac{p}{2}+6p)}, \\
&\quad 0, -2^{(-\frac{1}{2}+\frac{p}{2}+5p)}, -2^{5p}, 0, 0, -2^{(-\frac{1}{2}+\frac{p}{2}+3p)}, 0, \\
&\quad 2^{(-\frac{1}{2}+\frac{p}{2}+2p)}, 0, 0, 2^p, 2^{(-\frac{1}{2}+\frac{p}{2})}, 0)) , \tag{3.4}
\end{aligned}$$

$$\begin{aligned}
& \operatorname{Im} \operatorname{Li}_s \left[\frac{1}{\sqrt{2}^p} \exp \left(\frac{i\pi}{12} \right) \right] + \operatorname{Im} \operatorname{Li}_s \left[\frac{1}{\sqrt{2}^p} \exp \left(\frac{7i\pi}{12} \right) \right] \\
&= \frac{\sqrt{3}}{2^{12p}} P(s, 2^{12p}, 24, (2^{(-\frac{1}{2}+\frac{p}{2}+11p)}, 0, 0, 2^{10p}, \\
&\quad 2^{(-\frac{1}{2}+\frac{p}{2}+9p)}, 0, 2^{(-\frac{1}{2}+\frac{p}{2}+8p)}, 2^{8p}, 0, 0, 2^{(-\frac{1}{2}+\frac{p}{2}+6p)}, \\
&\quad 0, -2^{(-\frac{1}{2}+\frac{p}{2}+5p)}, 0, 0, -2^{4p}, -2^{(-\frac{1}{2}+\frac{p}{2}+3p)}, 0, \\
&\quad -2^{(-\frac{1}{2}+\frac{p}{2}+2p)}, -2^{2p}, 0, 0, -2^{(-\frac{1}{2}+\frac{p}{2})}, 0)) , \tag{3.5}
\end{aligned}$$

$$\begin{aligned}
& \operatorname{Im} \operatorname{Li}_s \left[\frac{1}{\sqrt{2^p}} \exp \left(\frac{i\pi}{12} \right) \right] - \operatorname{Im} \operatorname{Li}_s \left[\frac{1}{\sqrt{2^p}} \exp \left(\frac{7i\pi}{12} \right) \right] \\
&= \frac{1}{2^{12p}} P(s, 2^{12p}, 24, (-2^{(-\frac{1}{2}+\frac{p}{2}+11p)}, 2^{11p}, 2^{(\frac{1}{2}-\frac{p}{2}+11p)}, 0, \\
& 2^{(-\frac{1}{2}+\frac{p}{2}+9p)}, 2^{1+9p}, 2^{(-\frac{1}{2}+\frac{p}{2}+8p)}, 0, 2^{(\frac{1}{2}-\frac{p}{2}+8p)}, 2^{7p}, -2^{(-\frac{1}{2}+\frac{p}{2}+6p)}, \\
& 0, 2^{(-\frac{1}{2}+\frac{p}{2}+5p)}, -2^{5p}, -2^{(\frac{1}{2}-\frac{p}{2}+5p)}, 0, -2^{(-\frac{1}{2}+\frac{p}{2}+3p)}, -2^{1+3p}, \\
& -2^{(-\frac{1}{2}+\frac{p}{2}+2p)}, 0, -2^{(\frac{1}{2}-\frac{p}{2}+2p)}, -2^p, 2^{(-\frac{1}{2}+\frac{p}{2})}, 0)) , \tag{3.6}
\end{aligned}$$

$$\begin{aligned}
& \operatorname{Re} \operatorname{Li}_s \left[\frac{1}{\sqrt{2^p}} \exp \left(\frac{5i\pi}{12} \right) \right] + \operatorname{Re} \operatorname{Li}_s \left[\frac{1}{\sqrt{2^p}} \exp \left(\frac{11i\pi}{12} \right) \right] \\
&= \frac{1}{2^{12p}} P \left(s, 2^{12p}, 24, \left(-2^{(-\frac{1}{2}+\frac{p}{2}+11p)}, 0, -2^{(\frac{1}{2}-\frac{p}{2}+11p)}, 2^{10p}, \right. \right. \\
& 2^{(-\frac{1}{2}+\frac{p}{2}+9p)}, 0, -2^{(-\frac{1}{2}+\frac{p}{2}+8p)}, -2^{8p}, 2^{(\frac{1}{2}-\frac{p}{2}+8p)}, 0, 2^{(-\frac{1}{2}+\frac{p}{2}+6p)}, \\
& -2^{1+6p}, 2^{(-\frac{1}{2}+\frac{p}{2}+5p)}, 0, 2^{(\frac{1}{2}-\frac{p}{2}+5p)}, -2^{4p}, -2^{(-\frac{1}{2}+\frac{p}{2}+3p)}, 0, \\
& \left. \left. 2^{(-\frac{1}{2}+\frac{p}{2}+2p)}, 2^{2p}, -2^{(\frac{1}{2}-\frac{p}{2}+2p)}, 0, -2^{(-\frac{1}{2}+\frac{p}{2})}, 2 \right) \right) , \tag{3.7}
\end{aligned}$$

$$\begin{aligned}
& \operatorname{Re} \operatorname{Li}_s \left[\frac{1}{\sqrt{2^p}} \exp \left(\frac{5i\pi}{12} \right) \right] - \operatorname{Re} \operatorname{Li}_s \left[\frac{1}{\sqrt{2^p}} \exp \left(\frac{11i\pi}{12} \right) \right] \\
&= \frac{\sqrt{3}}{2^{12p}} P \left(s, 2^{12p}, 24, \left(2^{(-\frac{1}{2}+\frac{p}{2}+11p)}, -2^{11p}, 0, 0, \right. \right. \\
& 2^{(-\frac{1}{2}+\frac{p}{2}+9p)}, 0, -2^{(-\frac{1}{2}+\frac{p}{2}+8p)}, 0, 0, 2^{7p}, -2^{(-\frac{1}{2}+\frac{p}{2}+6p)}, \\
& 0, -2^{(-\frac{1}{2}+\frac{p}{2}+5p)}, 2^{5p}, 0, 0, -2^{(-\frac{1}{2}+\frac{p}{2}+3p)}, 0, \\
& \left. \left. 2^{(-\frac{1}{2}+\frac{p}{2}+2p)}, 0, 0, -2^p, 2^{(-\frac{1}{2}+\frac{p}{2})}, 0 \right) \right) , \tag{3.8}
\end{aligned}$$

$$\begin{aligned}
& \operatorname{Im} \operatorname{Li}_s \left[\frac{1}{\sqrt{2^p}} \exp \left(\frac{5i\pi}{12} \right) \right] + \operatorname{Im} \operatorname{Li}_s \left[\frac{1}{\sqrt{2^p}} \exp \left(\frac{11i\pi}{12} \right) \right] \\
&= \frac{\sqrt{3}}{2^{12p}} P \left(s, 2^{12p}, 24, \left(2^{(-\frac{1}{2} + \frac{p}{2} + 11p)}, 0, 0, -2^{10p}, \right. \right. \\
&\quad 2^{(-\frac{1}{2} + \frac{p}{2} + 9p)}, 0, 2^{(-\frac{1}{2} + \frac{p}{2} + 8p)}, -2^{8p}, 0, 0, 2^{(-\frac{1}{2} + \frac{p}{2} + 6p)}, \\
&\quad 0, -2^{(-\frac{1}{2} + \frac{p}{2} + 5p)}, 0, 0, 2^{4p}, -2^{(-\frac{1}{2} + \frac{p}{2} + 3p)}, 0, \\
&\quad \left. \left. -2^{(-\frac{1}{2} + \frac{p}{2} + 2p)}, 2^{2p}, 0, 0, -2^{(-\frac{1}{2} + \frac{p}{2})}, 0 \right) \right) \quad (3.9)
\end{aligned}$$

and

$$\begin{aligned}
& \operatorname{Im} \operatorname{Li}_s \left[\frac{1}{\sqrt{2^p}} \exp \left(\frac{5i\pi}{12} \right) \right] - \operatorname{Im} \operatorname{Li}_s \left[\frac{1}{\sqrt{2^p}} \exp \left(\frac{11i\pi}{12} \right) \right] \\
&= \frac{1}{2^{12p}} P \left(s, 2^{12p}, 24, \left(2^{(-\frac{1}{2} + \frac{p}{2} + 11p)}, 2^{11p}, -2^{(\frac{1}{2} - \frac{p}{2} + 11p)}, 0, \right. \right. \\
&\quad -2^{(-\frac{1}{2} + \frac{p}{2} + 9p)}, 2^{1+9p}, -2^{(-\frac{1}{2} + \frac{p}{2} + 8p)}, 0, -2^{(\frac{1}{2} - \frac{p}{2} + 8p)}, 2^{7p}, 2^{(-\frac{1}{2} + \frac{p}{2} + 6p)}, \\
&\quad 0, -2^{(-\frac{1}{2} + \frac{p}{2} + 5p)}, -2^{5p}, 2^{(\frac{1}{2} - \frac{p}{2} + 5p)}, 0, 2^{(-\frac{1}{2} + \frac{p}{2} + 3p)}, -2^{1+3p}, \\
&\quad \left. \left. 2^{(-\frac{1}{2} + \frac{p}{2} + 2p)}, 0, 2^{(\frac{1}{2} - \frac{p}{2} + 2p)}, -2^p, -2^{(-\frac{1}{2} + \frac{p}{2})}, 0 \right) \right) \quad (3.10)
\end{aligned}$$

We note that although the above formulas hold true for all $p > 0$, they are BBP-type only for $p \in \mathbb{Z}^+$ and $\bmod(p, 2) = 1$.

4 Degree 1 Formulas

When $s = 1$, the polylogarithm constants on the left hand side in each of the above formulas can be evaluated, using the identities

$$\arctan \left(\frac{q \sin x}{1 - q \cos x} \right) = \operatorname{Im} \operatorname{Li}_1 [q \exp(ix)]$$

and

$$-\frac{1}{2} \log (1 - 2q \cos x + q^2) = \operatorname{Re} \operatorname{Li}_1 [q \exp(ix)] ,$$

and we have the following degree 1 binary BBP-type formulas

$$\begin{aligned}
& -\frac{1}{2} \log(1 - 2^{(\frac{1}{2}-\frac{p}{2})} + 2^{-p} - 2^{(\frac{1}{2}+\frac{p}{2}-2p)} + 2^{-2p}) \\
& = \frac{1}{2^{12p}} P(1, 2^{12p}, 24, (2^{(-\frac{1}{2}+\frac{p}{2}+11p)}, 0, 2^{(\frac{1}{2}-\frac{p}{2}+11p)}, 2^{10p}, \\
& \quad -2^{(-\frac{1}{2}+\frac{p}{2}+9p)}, 0, 2^{(-\frac{1}{2}+\frac{p}{2}+8p)}, -2^{8p}, -2^{(\frac{1}{2}-\frac{p}{2}+8p)}, 0, -2^{(-\frac{1}{2}+\frac{p}{2}+6p)}, \\
& \quad -2^{1+6p}, -2^{(-\frac{1}{2}+\frac{p}{2}+5p)}, 0, -2^{(\frac{1}{2}-\frac{p}{2}+5p)}, -2^{4p}, 2^{(-\frac{1}{2}+\frac{p}{2}+3p)}, 0, \\
& \quad -2^{(-\frac{1}{2}+\frac{p}{2}+2p)}, 2^{2p}, 2^{(\frac{1}{2}-\frac{p}{2}+2p)}, 0, 2^{(-\frac{1}{2}+\frac{p}{2})}, 2)), \tag{4.1}
\end{aligned}$$

$$\begin{aligned}
& \frac{\sqrt{3}}{2} \log \left(\frac{(1 + 2^{(-\frac{1}{2}-\frac{p}{2})}) \sqrt{3} - 2^{(-\frac{1}{2}-\frac{p}{2})} + 2^{-p})^2}{1 - 2^{(\frac{1}{2}-\frac{p}{2})} - 2^{(\frac{1}{2}-\frac{3p}{2})} + 2^{-2p} + 2^{-p}} \right) \\
& = \frac{3}{2^{12p}} P(1, 2^{12p}, 24, (2^{(-\frac{1}{2}+\frac{p}{2}+11p)}, 2^{11p}, 0, 0, \\
& \quad 2^{(-\frac{1}{2}+\frac{p}{2}+9p)}, 0, -2^{(-\frac{1}{2}+\frac{p}{2}+8p)}, 0, 0, -2^{7p}, \\
& \quad -2^{(-\frac{1}{2}+\frac{p}{2}+6p)}, 0, -2^{(-\frac{1}{2}+\frac{p}{2}+5p)}, -2^{5p}, 0, 0, \\
& \quad -2^{(-\frac{1}{2}+\frac{p}{2}+3p)}, 0, 2^{(-\frac{1}{2}+\frac{p}{2}+2p)}, 0, 0, 2^p, \\
& \quad 2^{(-\frac{1}{2}+\frac{p}{2})}, 0)), \tag{4.2}
\end{aligned}$$

$$\begin{aligned}
& \sqrt{3} \arctan \left[\left(\frac{1 - 2^{\frac{p+1}{2}}}{1 + 2^{\frac{p+1}{2}} - 2^{p+1}} \right) \sqrt{3} \right] \\
& = \frac{3}{2^{12p}} P(1, 2^{12p}, 24, (2^{(-\frac{1}{2}+\frac{p}{2}+11p)}, \\
& \quad 0, 0, 2^{10p}, 2^{(-\frac{1}{2}+\frac{p}{2}+9p)}, 0, 2^{(-\frac{1}{2}+\frac{p}{2}+8p)}, 2^{8p}, 0, 0, 2^{(-\frac{1}{2}+\frac{p}{2}+6p)}, \\
& \quad 0, -2^{(-\frac{1}{2}+\frac{p}{2}+5p)}, 0, 0, -2^{4p}, -2^{(-\frac{1}{2}+\frac{p}{2}+3p)}, 0, \\
& \quad -2^{(-\frac{1}{2}+\frac{p}{2}+2p)}, -2^{2p}, 0, 0, -2^{(-\frac{1}{2}+\frac{p}{2})}, 0)), \tag{4.3}
\end{aligned}$$

$$\begin{aligned}
-\arctan \left[\frac{2^{\frac{1-p}{2}} - 1}{-2^{\frac{1+p}{2}} + 1} \right] &= \frac{1}{2^{12p}} P(1, 2^{12p}, 24, (-2^{(-\frac{1}{2}+\frac{p}{2}+11p)}, 2^{11p}, 2^{(\frac{1}{2}-\frac{p}{2}+11p)}, \\
&0, 2^{(-\frac{1}{2}+\frac{p}{2}+9p)}, 2^{1+9p}, 2^{(-\frac{1}{2}+\frac{p}{2}+8p)}, 0, 2^{(\frac{1}{2}-\frac{p}{2}+8p)}, 2^{7p}, -2^{(-\frac{1}{2}+\frac{p}{2}+6p)}, \\
&0, 2^{(-\frac{1}{2}+\frac{p}{2}+5p)}, -2^{5p}, -2^{(\frac{1}{2}-\frac{p}{2}+5p)}, 0, -2^{(-\frac{1}{2}+\frac{p}{2}+3p)}, -2^{1+3p}, \\
&-2^{(-\frac{1}{2}+\frac{p}{2}+2p)}, 0, -2^{(\frac{1}{2}-\frac{p}{2}+2p)}, -2^p, 2^{(-\frac{1}{2}+\frac{p}{2})}, 0)), \quad (4.4)
\end{aligned}$$

$$\begin{aligned}
&-\frac{1}{2} \log(1 + 2^{(\frac{1}{2}-\frac{p}{2})} + 2^{-p} + 2^{(\frac{1}{2}+\frac{p}{2}-2p)} + 2^{-2p}) \\
&= \frac{1}{2^{12p}} P \left(1, 2^{12p}, 24, \left(-2^{(-\frac{1}{2}+\frac{p}{2}+11p)}, 0, -2^{(\frac{1}{2}-\frac{p}{2}+11p)}, 2^{10p}, \right. \right. \\
&2^{(-\frac{1}{2}+\frac{p}{2}+9p)}, 0, -2^{(-\frac{1}{2}+\frac{p}{2}+8p)}, -2^{8p}, 2^{(\frac{1}{2}-\frac{p}{2}+8p)}, 0, 2^{(-\frac{1}{2}+\frac{p}{2}+6p)}, \\
&-2^{1+6p}, 2^{(-\frac{1}{2}+\frac{p}{2}+5p)}, 0, 2^{(\frac{1}{2}-\frac{p}{2}+5p)}, -2^{4p}, -2^{(-\frac{1}{2}+\frac{p}{2}+3p)}, 0, \\
&\left. \left. 2^{(-\frac{1}{2}+\frac{p}{2}+2p)}, 2^{2p}, -2^{(\frac{1}{2}-\frac{p}{2}+2p)}, 0, -2^{(-\frac{1}{2}+\frac{p}{2})}, 2 \right) \right), \quad (4.5)
\end{aligned}$$

$$\begin{aligned}
\frac{\sqrt{3}}{2} \log \left[\frac{1 + 2^{-\frac{1}{2}-\frac{p}{2}}(1 + \sqrt{3}) + 2^{-p}}{1 + 2^{-\frac{1}{2}-\frac{p}{2}}(1 - \sqrt{3}) + 2^{-p}} \right] &= \frac{3}{2^{12p}} P \left(1, 2^{12p}, 24, \left(2^{(-\frac{1}{2}+\frac{p}{2}+11p)}, -2^{11p}, 0, 0, \right. \right. \\
&2^{(-\frac{1}{2}+\frac{p}{2}+9p)}, 0, -2^{(-\frac{1}{2}+\frac{p}{2}+8p)}, 0, 0, 2^{7p}, -2^{(-\frac{1}{2}+\frac{p}{2}+6p)}, \\
&0, -2^{(-\frac{1}{2}+\frac{p}{2}+5p)}, 2^{5p}, 0, 0, -2^{(-\frac{1}{2}+\frac{p}{2}+3p)}, 0, \\
&\left. \left. 2^{(-\frac{1}{2}+\frac{p}{2}+2p)}, 0, 0, -2^p, 2^{(-\frac{1}{2}+\frac{p}{2})}, 0 \right) \right), \quad (4.6)
\end{aligned}$$

$$\begin{aligned}
\sqrt{3} \arctan \left[\left(\frac{1 + 2^{\frac{p+1}{2}}}{-1 + 2^{\frac{p+1}{2}} + 2^{p+1}} \right) \sqrt{3} \right] &= \frac{3}{2^{12p}} P \left(1, 2^{12p}, 24, \left(2^{(-\frac{1}{2}+\frac{p}{2}+11p)}, 0, 0, -2^{10p}, \right. \right. \\
&2^{(-\frac{1}{2}+\frac{p}{2}+9p)}, 0, 2^{(-\frac{1}{2}+\frac{p}{2}+8p)}, -2^{8p}, 0, 0, 2^{(-\frac{1}{2}+\frac{p}{2}+6p)}, \\
&0, -2^{(-\frac{1}{2}+\frac{p}{2}+5p)}, 0, 0, 2^{4p}, -2^{(-\frac{1}{2}+\frac{p}{2}+3p)}, 0, \\
&\left. \left. -2^{(-\frac{1}{2}+\frac{p}{2}+2p)}, 2^{2p}, 0, 0, -2^{(-\frac{1}{2}+\frac{p}{2})}, 0 \right) \right), \quad (4.7)
\end{aligned}$$

and

$$\begin{aligned} \arctan \left[\frac{2^{\frac{1-p}{2}} + 1}{2^{\frac{1+p}{2}} + 1} \right] &= \frac{1}{2^{12p}} P \left(1, 2^{12p}, 24, \left(2^{(-\frac{1}{2} + \frac{p}{2} + 11p)}, 2^{11p}, -2^{(\frac{1}{2} - \frac{p}{2} + 11p)}, 0, \right. \right. \\ &\quad \left. \left. -2^{(-\frac{1}{2} + \frac{p}{2} + 9p)}, 2^{1+9p}, -2^{(-\frac{1}{2} + \frac{p}{2} + 8p)}, 0, -2^{(\frac{1}{2} - \frac{p}{2} + 8p)}, 2^{7p}, 2^{(-\frac{1}{2} + \frac{p}{2} + 6p)}, \right. \right. \\ &\quad \left. \left. 0, -2^{(-\frac{1}{2} + \frac{p}{2} + 5p)}, -2^{5p}, 2^{(\frac{1}{2} - \frac{p}{2} + 5p)}, 0, 2^{(-\frac{1}{2} + \frac{p}{2} + 3p)}, -2^{1+3p}, \right. \right. \\ &\quad \left. \left. 2^{(-\frac{1}{2} + \frac{p}{2} + 2p)}, 0, 2^{(\frac{1}{2} - \frac{p}{2} + 2p)}, -2^p, -2^{(-\frac{1}{2} + \frac{p}{2})}, 0 \right) \right). \end{aligned} \quad (4.8)$$

Particular cases of these formulas will be discussed in section 6.1.

5 Degree 2 Formulas

The imaginary part of the dilogarithm function can be expressed in closed form as [6]

$$\operatorname{Im} \operatorname{Li}_2 [qe^{ix}] = \omega \log q + \frac{1}{2} \operatorname{Cl}_2(2\omega) - \frac{1}{2} \operatorname{Cl}_2(2\omega + 2x) + \frac{1}{2} \operatorname{Cl}_2(2x), \quad (5.1)$$

where

$$\omega = \arctan \left(\frac{q \sin x}{1 - q \cos x} \right).$$

Using (5.1), the results (3.5) and (3.6) can be written in degree 2 as

$$\begin{aligned} &-(\omega_1 + \omega_2)p \log 2 + \operatorname{Cl}_2(2\omega_1) + \operatorname{Cl}_2(2\omega_2) \\ &-\operatorname{Cl}_2(2\omega_1 + \pi/6) + \operatorname{Cl}_2(-2\omega_2 + 5\pi/6) + 1/2 \operatorname{Cl}_2(\pi/3) \\ &= 2 \operatorname{Im} \operatorname{Li}_2 \left[\frac{1}{\sqrt{2^p}} \exp \left(\frac{i\pi}{12} \right) \right] + 2 \operatorname{Im} \operatorname{Li}_2 \left[\frac{1}{\sqrt{2^p}} \exp \left(\frac{7i\pi}{12} \right) \right] \\ &= \frac{\sqrt{3}}{2^{12p-1}} P(2, 2^{12p}, 24, (2^{(-\frac{1}{2} + \frac{p}{2} + 11p)}, 0, 0, 2^{10p}, \\ &\quad 2^{(-\frac{1}{2} + \frac{p}{2} + 9p)}, 0, 2^{(-\frac{1}{2} + \frac{p}{2} + 8p)}, 2^{8p}, 0, 0, 2^{(-\frac{1}{2} + \frac{p}{2} + 6p)}, \\ &\quad 0, -2^{(-\frac{1}{2} + \frac{p}{2} + 5p)}, 0, 0, -2^{4p}, -2^{(-\frac{1}{2} + \frac{p}{2} + 3p)}, 0, \\ &\quad -2^{(-\frac{1}{2} + \frac{p}{2} + 2p)}, -2^{2p}, 0, 0, -2^{(-\frac{1}{2} + \frac{p}{2})}, 0)) \end{aligned} \quad (5.2)$$

and

$$\begin{aligned}
& (\omega_2 - \omega_1)p \log 2 + \text{Cl}_2(2\omega_1) - \text{Cl}_2(2\omega_2) \\
& - \text{Cl}_2(2\omega_1 + \pi/6) - \text{Cl}_2(-2\omega_2 + 5\pi/6) + 4G/3 \\
& = 2 \text{Im Li}_2 \left[\frac{1}{\sqrt{2}^p} \exp \left(\frac{i\pi}{12} \right) \right] - 2 \text{Im Li}_2 \left[\frac{1}{\sqrt{2}^p} \exp \left(\frac{7i\pi}{12} \right) \right] \\
& = \frac{1}{2^{12p-1}} P(2, 2^{12p}, 24, (-2^{(-\frac{1}{2}+\frac{p}{2}+11p)}, 2^{11p}, 2^{(\frac{1}{2}-\frac{p}{2}+11p)}, 0, \\
& 2^{(-\frac{1}{2}+\frac{p}{2}+9p)}, 2^{1+9p}, 2^{(-\frac{1}{2}+\frac{p}{2}+8p)}, 0, 2^{(\frac{1}{2}-\frac{p}{2}+8p)}, 2^{7p}, -2^{(-\frac{1}{2}+\frac{p}{2}+6p)}, \\
& 0, 2^{(-\frac{1}{2}+\frac{p}{2}+5p)}, -2^{5p}, -2^{(\frac{1}{2}-\frac{p}{2}+5p)}, 0, -2^{(-\frac{1}{2}+\frac{p}{2}+3p)}, -2^{1+3p}, \\
& -2^{(-\frac{1}{2}+\frac{p}{2}+2p)}, 0, -2^{(\frac{1}{2}-\frac{p}{2}+2p)}, -2^p, 2^{(-\frac{1}{2}+\frac{p}{2})}, 0)) , \tag{5.3}
\end{aligned}$$

where ω_1 and ω_2 are given by

$$\tan \omega_1 = \frac{\sqrt{3} - 1}{\sqrt{2}^{p+3} - \sqrt{3} - 1}$$

and

$$\tan \omega_2 = \frac{\sqrt{3} + 1}{\sqrt{2}^{p+3} + \sqrt{3} - 1}.$$

Similarly, using (5.1), the results (3.9) and (3.10) can be written in degree 2 as

$$\begin{aligned}
& -(\omega_3 + \omega_4)p \log 2 + \text{Cl}_2(2\omega_3) + \text{Cl}_2(2\omega_4) \\
& - \text{Cl}_2(2\omega_3 + 5\pi/6) + \text{Cl}_2(-2\omega_4 + \pi/6) - 1/2 \text{Cl}_2(\pi/3) \\
& = 2 \text{Im Li}_2 \left[\frac{1}{\sqrt{2}^p} \exp \left(\frac{5i\pi}{12} \right) \right] + 2 \text{Im Li}_2 \left[\frac{1}{\sqrt{2}^p} \exp \left(\frac{11i\pi}{12} \right) \right] \\
& = \frac{\sqrt{3}}{2^{12p-1}} P \left(2, 2^{12p}, 24, \left(2^{(-\frac{1}{2}+\frac{p}{2}+11p)}, 0, 0, -2^{10p}, \right. \right. \\
& 2^{(-\frac{1}{2}+\frac{p}{2}+9p)}, 0, 2^{(-\frac{1}{2}+\frac{p}{2}+8p)}, -2^{8p}, 0, 0, 2^{(-\frac{1}{2}+\frac{p}{2}+6p)}, \\
& 0, -2^{(-\frac{1}{2}+\frac{p}{2}+5p)}, 0, 0, 2^{4p}, -2^{(-\frac{1}{2}+\frac{p}{2}+3p)}, 0, \\
& \left. \left. -2^{(-\frac{1}{2}+\frac{p}{2}+2p)}, 2^{2p}, 0, 0, -2^{(-\frac{1}{2}+\frac{p}{2})}, 0 \right) \right) \tag{5.4}
\end{aligned}$$

and

$$\begin{aligned}
& (\omega_4 - \omega_3)p \log 2 + \text{Cl}_2(2\omega_3) - \text{Cl}_2(2\omega_4) \\
& - \text{Cl}_2(2\omega_3 + 5\pi/6) - \text{Cl}_2(-2\omega_4 + \pi/6) + 4G/3 \\
& = 2 \text{Im Li}_2 \left[\frac{1}{\sqrt{2}^p} \exp \left(\frac{5i\pi}{12} \right) \right] - 2 \text{Im Li}_2 \left[\frac{1}{\sqrt{2}^p} \exp \left(\frac{11i\pi}{12} \right) \right] \\
& = \frac{1}{2^{12p-1}} P \left(2, 2^{12p}, 24, \left(2^{(-\frac{1}{2} + \frac{p}{2} + 11p)}, 2^{11p}, -2^{(\frac{1}{2} - \frac{p}{2} + 11p)}, 0, \right. \right. \\
& \quad \left. \left. -2^{(-\frac{1}{2} + \frac{p}{2} + 9p)}, 2^{1+9p}, -2^{(-\frac{1}{2} + \frac{p}{2} + 8p)}, 0, -2^{(\frac{1}{2} - \frac{p}{2} + 8p)}, 2^{7p}, 2^{(-\frac{1}{2} + \frac{p}{2} + 6p)}, \right. \right. \\
& \quad \left. \left. 0, -2^{(-\frac{1}{2} + \frac{p}{2} + 5p)}, -2^{5p}, 2^{(\frac{1}{2} - \frac{p}{2} + 5p)}, 0, 2^{(-\frac{1}{2} + \frac{p}{2} + 3p)}, -2^{1+3p}, \right. \right. \\
& \quad \left. \left. 2^{(-\frac{1}{2} + \frac{p}{2} + 2p)}, 0, 2^{(\frac{1}{2} - \frac{p}{2} + 2p)}, -2^p, -2^{(-\frac{1}{2} + \frac{p}{2})}, 0 \right) \right), \tag{5.5}
\end{aligned}$$

where ω_3 and ω_4 are given by

$$\tan \omega_3 = \frac{\sqrt{3} + 1}{\sqrt{2}^{p+3} - \sqrt{3} + 1}$$

and

$$\tan \omega_4 = \frac{\sqrt{3} - 1}{\sqrt{2}^{p+3} + \sqrt{3} + 1}.$$

In the above formulas, $G = \text{Cl}_2(\pi/2)$ is Catalan's constant.

In deriving (5.4) and (5.5), we used equation 4.32 pg 106 and equation 4.17 pg 104 of [6], namely,

$$\text{Cl}_2 \left(\frac{\pi}{6} \right) + \text{Cl}_2 \left(\frac{5\pi}{6} \right) = \frac{4G}{3}$$

and

$$\text{Cl}_2(x) - \text{Cl}_2(\pi - x) = \frac{1}{2} \text{Cl}_2(2x).$$

6 Interesting Particular Cases

6.1 Degree 1 Binary BBP-type Formulas

We first note that (4.1) and (4.5) give an infinite set of primes whose logarithms have binary BBP-type formulas, which serve to augment the known

ones (e.g. those found in [2]) and [5]. Similarly, (4.4) and (4.8) give an infinite set of rationals whose arctangents have binary BBP-type representations. We now present a couple of binary degree 1 formulas.

6.1.1 Binary BBP-type formula for $\log 2$

The identity

$$\log 2 = \operatorname{Re} \operatorname{Li}_1 \left[\frac{1}{\sqrt{2}} \exp \left(\frac{i\pi}{12} \right) \right] + \operatorname{Re} \operatorname{Li}_1 \left[\frac{1}{\sqrt{2}} \exp \left(\frac{7i\pi}{12} \right) \right] \quad (6.1)$$

and $p = 1$ in (4.1) lead to the binary BBP-type formula

$$\begin{aligned} \log 2 = & \frac{1}{2^{12}} P(1, 2^{12}, 24, (2^{11}, 0, 2^{11}, 2^{10}, -2^9, 0, 2^8, -2^8, \\ & -2^8, 0, -2^6, -2^7, -2^5, 0, -2^5, -2^4, 2^3, 0, \\ & -2^2, 2^2, 2^2, 0, 1, 2)) . \end{aligned} \quad (6.2)$$

6.1.2 Binary BBP-type formula for $\pi\sqrt{3}$

The identity

$$\frac{\pi}{3} = \operatorname{Im} \operatorname{Li}_1 \left[\frac{1}{\sqrt{2}} \exp \left(\frac{i\pi}{12} \right) \right] + \operatorname{Im} \operatorname{Li}_1 \left[\frac{1}{\sqrt{2}} \exp \left(\frac{7i\pi}{12} \right) \right] \quad (6.3)$$

and $p = 1$ in (4.3) lead to the binary BBP-type formula

$$\begin{aligned} \pi\sqrt{3} = & \frac{9}{2^{12}} P(1, 2^{12}, 24, (2^{11}, 0, 0, 2^{10}, 2^9, 0, 2^8, 2^8, 0, 0, \\ & 2^6, 0, -2^5, 0, 0, -2^4, -2^3, 0, -2^2, -2^2, 0, 0, -1, 0)) . \end{aligned} \quad (6.4)$$

6.1.3 Binary BBP-type formula for $\sqrt{3}\log(2 + \sqrt{3})$

The identity

$$\log(2 + \sqrt{3}) = \operatorname{Re} \operatorname{Li}_1 \left[\frac{1}{\sqrt{2}} \exp \left(\frac{i\pi}{12} \right) \right] - \operatorname{Re} \operatorname{Li}_1 \left[\frac{1}{\sqrt{2}} \exp \left(\frac{7i\pi}{12} \right) \right] \quad (6.5)$$

and $p = 1$ in (4.2) lead to the binary BBP-type formula

$$\begin{aligned} \sqrt{3} \log(2 + \sqrt{3}) = \frac{3}{2^{12}} P(1, 2^{12}, 24, (2^{11}, 2^{11}, 0, 0, 2^9, 0, -2^8, \\ 0, 0, -2^7, -2^6, 0, -2^5, -2^5, 0, 0, -2^3, \\ 0, 2^2, 0, 0, 2, 1, 0)) \end{aligned} \quad (6.6)$$

6.1.4 Binary BBP-type formula for $\arctan 1/6$

The identity

$$-\arctan\left(\frac{1}{6}\right) = \operatorname{Im} \operatorname{Li}_1 \left[\frac{1}{\sqrt{2}^3} \exp\left(\frac{i\pi}{12}\right) \right] - \operatorname{Im} \operatorname{Li}_1 \left[\frac{1}{\sqrt{2}^3} \exp\left(\frac{7i\pi}{12}\right) \right] \quad (6.7)$$

and $p = 3$ in (4.4) lead to the binary BBP-type formula

$$\begin{aligned} \arctan\left(\frac{1}{6}\right) = \frac{1}{2^{35}} P(1, 2^{36}, 24, (2^{33}, -2^{32}, -2^{31}, 0, -2^{27}, \\ -2^{27}, -2^{24}, 0, -2^{22}, -2^{20}, 2^{18}, 0, -2^{15}, 2^{14}, \\ 2^{13}, 0, 2^9, 2^9, 2^6, 0, 2^4, 2^2, -1, 0)) . \end{aligned} \quad (6.8)$$

6.2 Degree 2 Binary BBP-type Formulas

When $p = 1$ then $\omega_1 = \omega_2 = \pi/6$ and (5.4) and (5.5) simplify to

$$-\frac{\pi}{3} \log 2 + \frac{5}{2} \operatorname{Cl}_2\left(\frac{\pi}{3}\right) = 2 \operatorname{Im} \operatorname{Li}_2 \left[\frac{1}{\sqrt{2}} \exp\left(\frac{i\pi}{12}\right) \right] + 2 \operatorname{Im} \operatorname{Li}_2 \left[\frac{1}{\sqrt{2}} \exp\left(\frac{7i\pi}{12}\right) \right] \quad (6.9)$$

and

$$G = 3 \operatorname{Im} \operatorname{Li}_2 \left[\frac{1}{\sqrt{2}} \exp\left(\frac{i7\pi}{12}\right) \right] - 3 \operatorname{Im} \operatorname{Li}_2 \left[\frac{1}{\sqrt{2}} \exp\left(\frac{i\pi}{12}\right) \right] , \quad (6.10)$$

respectively.

Choosing $q = 1/2$ and $x = \pi/3$ in (5.1) gives

$$-\pi \log 2 + 5 \operatorname{Cl}_2\left(\frac{\pi}{3}\right) = 6 \operatorname{Im} \operatorname{Li}_2 \left[\frac{1}{2} \exp\left(\frac{\pi}{3}\right) \right] . \quad (6.11)$$

We note that

$$\begin{aligned} \operatorname{Im} \operatorname{Li}_2 \left[\frac{1}{2} \exp \left(\frac{i\pi}{3} \right) \right] &= \sum_{k=1}^{\infty} \frac{1}{2^k} \frac{\sin(k\pi/3)}{k^2} \\ &= \frac{\sqrt{3}}{2^{10}} P(2, 2^{12}, 24, (0, 2^{10}, 0, 2^9, 0, 0, 0, -2^7, 0, -2^6, 0, 0, 0, \\ &\quad 2^4, 0, 2^3, 0, 0, 0, -2, 0, -1, 0, 0)). \end{aligned} \quad (6.12)$$

6.2.1 Binary BBP-type formula for $\sqrt{3} \operatorname{Cl}_2(\pi/3)$

Eliminating $\pi \log 2$ between (6.9) and (6.11) we have

$$\begin{aligned} \operatorname{Cl}_2 \left(\frac{\pi}{3} \right) &= \frac{12}{5} \left\{ \operatorname{Im} \operatorname{Li}_2 \left[\frac{1}{\sqrt{2}} \exp \left(\frac{i\pi}{12} \right) \right] + \operatorname{Im} \operatorname{Li}_2 \left[\frac{1}{\sqrt{2}} \exp \left(\frac{7i\pi}{12} \right) \right] \right\} \\ &\quad - \frac{12}{5} \operatorname{Im} \operatorname{Li}_2 \left[\frac{1}{2} \exp \left(\frac{i\pi}{3} \right) \right]. \end{aligned} \quad (6.13)$$

Using (5.4) (with $p = 1$) and (6.12) in (6.13), we obtain a binary BBP-type formula for $\sqrt{3} \operatorname{Cl}_2(\pi/3)$:

$$\begin{aligned} \sqrt{3} \operatorname{Cl}_2 \left(\frac{\pi}{3} \right) &= \frac{9}{5 \cdot 2^{10}} P(2, 2^{12}, 24, (2^{11}, -2^{12}, 0, -2^{10}, 2^9, \\ &\quad 0, 2^8, 3 \cdot 2^8, 0, 2^8, 2^6, 0, -2^5, -2^6, 0, \\ &\quad -3 \cdot 2^4, -2^3, 0, -2^2, 2^2, 0, 2^2, -1, 0)). \end{aligned} \quad (6.14)$$

6.2.2 Binary BBP-type formula for $\pi\sqrt{3} \log 2$

Eliminating $\operatorname{Cl}_2(\pi/3)$ between (6.9) and (6.11) we have

$$\begin{aligned} \pi \log 2 &= 12 \left\{ \operatorname{Im} \operatorname{Li}_2 \left[\frac{1}{\sqrt{2}} \exp \left(\frac{i\pi}{12} \right) \right] + \operatorname{Im} \operatorname{Li}_2 \left[\frac{1}{\sqrt{2}} \exp \left(\frac{7i\pi}{12} \right) \right] \right\} \\ &\quad - 18 \operatorname{Im} \operatorname{Li}_2 \left[\frac{1}{2} \exp \left(\frac{i\pi}{3} \right) \right]. \end{aligned} \quad (6.15)$$

Using (5.4) (with $p = 1$) and (6.12) in (6.15), we obtain a binary BBP-type formula for $\pi\sqrt{3} \log 2$:

$$\begin{aligned} \pi\sqrt{3} \log 2 &= \frac{9}{2^{10}} P(2, 2^{12}, 24, (2^{11}, -3 \cdot 2^{11}, 0, -2^{11}, 2^9, 0, 2^8, 2^{10}, 0, 3 \cdot 2^7, 2^6, 0, \\ &\quad -2^5, -3 \cdot 2^5, 0, -2^6, -2^3, 0, -2^2, 2^3, 0, 6, -1, 0)). \end{aligned} \quad (6.16)$$

A variant of formula (6.16) (formula 32, section 5, in the BBP-Compendium) was discovered experimentally over ten years ago, but is hitherto unproved. Formula 32 in the Compendium is now proved by adding rational multiples of two zero relations to (6.16) (See [2]).

6.2.3 Binary BBP-type formula for Catalan's constant G

(6.10) leads immediately to

$$G = \frac{3}{2^{12}} P(2, 2^{12}, 24, (2^{11}, -2^{11}, -2^{11}, 0, -2^9, -2^{10}, -2^8, 0, -2^8, -2^7, 2^6, 0, -2^5, 2^5, 2^5, 0, 2^3, 2^4, 2^2, 0, 2^2, 2, 1, 0)) . \quad (6.17)$$

6.3 Binary Zero Relations

The identity

$$\frac{\pi}{6} = \text{Im Li}_1 \left[\frac{1}{2} \exp \left(\frac{i\pi}{3} \right) \right] = \sum_{k=1}^{\infty} \frac{1}{2^k} \frac{\sin(k\pi/3)}{k} \quad (6.18)$$

leads to the binary BBP-type formula

$$\pi\sqrt{3} = \frac{9}{2^{10}} P(1, 2^{12}, 24, (0, 2^{10}, 0, 2^9, 0, 0, 0, -2^7, 0, -2^6, 0, 0, 0, 2^4, 0, 2^3, 0, 0, 0, -2, 0, -1, 0, 0)) . \quad (6.19)$$

We note that this is the base 2^{12} version of the formula listed for $\pi\sqrt{3}$ in section 4 of the BBP Compendium [2].

Combining (6.3) and (6.18), we have the identity

$$0 = \text{Im Li}_1 \left[\frac{1}{\sqrt{2}} \exp \left(\frac{i\pi}{12} \right) \right] + \text{Im Li}_1 \left[\frac{1}{\sqrt{2}} \exp \left(\frac{7i\pi}{12} \right) \right] - 2 \text{Im Li}_1 \left[\frac{1}{2} \exp \left(\frac{i\pi}{3} \right) \right] , \quad (6.20)$$

which, with the use of (6.4) and (6.19) gives the binary zero relation

$$0 = P(1, 2^{12}, 24, (2^{11}, -2^{12}, 0, -2^{10}, 2^9, 0, 2^8, 3 \cdot 2^8, 0, 2^8, 2^6, 0, -2^5, -2^6, 0, -3 \cdot 2^4, -2^3, 0, -2^2, 2^2, 0, 2^2, -1, 0)) . \quad (6.21)$$

(6.1) and the formula $\text{Li}_1(1/2) = \log 2$ give the identity

$$0 = \text{Re Li}_1 \left[\frac{1}{\sqrt{2}} \exp \left(\frac{i\pi}{12} \right) \right] + \text{Re Li}_1 \left[\frac{1}{\sqrt{2}} \exp \left(\frac{7i\pi}{12} \right) \right] - \text{Li}_1 \left[\frac{1}{2} \right] \quad (6.22)$$

which leads to the binary BBP-type zero relation

$$0 = P(1, 2^{12}, 24, (2^{11}, -2^{12}, 2^{11}, -2^{10}, -2^9, -2^{10}, 2^8, -3 \cdot 2^8, -2^8, -2^8, -2^6, -2^8, -2^5, -2^6, -2^5, -3 \cdot 2^4, 2^3, -2^4, -2^2, -2^2, 2^2, -2^2, 1, 0)). \quad (6.23)$$

The identity

$$0 = \text{Im Li}_1 \left[\frac{1}{\sqrt{2}} \exp \left(\frac{i\pi}{12} \right) \right] - \text{Im Li}_1 \left[\frac{1}{\sqrt{2}} \exp \left(\frac{7i\pi}{12} \right) \right] \quad (6.24)$$

and $p = 1$ in (4.4) lead to the binary BBP-type zero relation

$$0 = P(1, 2^{12}, 24, (-2^{11}, 2^{11}, 2^{11}, 0, 2^9, 2^{10}, 2^8, 0, 2^8, 2^7, -2^6, 0, 2^5, -2^5, -2^5, 0, -2^3, -2^4, -2^2, 0, -2^2, -2, 1, 0)). \quad (6.25)$$

The identity

$$0 = \text{Re Li}_1 \left[\frac{1}{\sqrt{2}} \exp \left(\frac{i\pi}{12} \right) \right] + \text{Re Li}_1 \left[\frac{1}{\sqrt{2}} \exp \left(\frac{7i\pi}{12} \right) \right] - 2 \text{Re Li}_1 \left[\frac{1}{\sqrt{2}} \exp \left(\frac{i\pi}{4} \right) \right] \quad (6.26)$$

leads to the binary zero relation

$$0 = P(1, 2^{12}, 24, (2^{11}, 0, -2^{12}, -3 \cdot 2^{10}, -2^9, 0, 2^8, 3 \cdot 2^8, 2^9, 0, -2^6, 0, -2^5, 0, 2^6, 3 \cdot 2^4, 2^3, 0, -2^2, -3 \cdot 2^2, -2^3, 0, 1, 0)). \quad (6.27)$$

The identity

$$0 = \text{Re Li}_1 \left[\frac{1}{\sqrt{2}} \exp \left(\frac{i\pi}{12} \right) \right] + \text{Re Li}_1 \left[\frac{1}{\sqrt{2}} \exp \left(\frac{7i\pi}{12} \right) \right] + 2 \text{Re Li}_1 \left[\frac{1}{\sqrt{2}} \exp \left(\frac{i\pi}{2} \right) \right] - 2 \text{Re Li}_1 \left[\frac{1}{2} \exp \left(\frac{i\pi}{3} \right) \right] \quad (6.28)$$

leads to the binary zero relation

$$0 = P(1, 2^{12}, 24, (2^{11}, -2^{13}, 2^{11}, 5 \cdot 2^{10}, -2^9, 2^{10}, 2^8, 3 \cdot 2^8, -2^8, -2^9, -2^6, -2^8, -2^5, -2^7, -2^5, 3 \cdot 2^4, 2^3, 2^4, -2^2, 5 \cdot 2^2, 2^2, -2^3, 1, 0)). \quad (6.29)$$

More degree 1 binary BBP-type zero relations in base 2^{12} and other bases can also be found in [7].

7 Conclusion

Using a clear and straightforward approach, explicit digit extraction BBP-type formulas in very general binary bases were discovered. As particular examples, new binary formulas were obtained for $\pi\sqrt{3}$, $\sqrt{3}\pi \log 2$ and some other polylogarithm constants. New binary BBP-type zero relations were also established.

References

- [1] David H. Bailey, Peter B. Borwein, and Simon Plouffe. On the rapid computation of various polylogarithmic constants. *Mathematics of Computation*, 66(218):903–913, 1997.
- [2] David H. Bailey. A compendium of BBP-type formulas for mathematical constants
<http://crd.lbl.gov/~dhbailey/dhbpapers/bbp-formulas.pdf>.
April 2013.
- [3] David H. Bailey and Richard E. Crandall. On the random character of fundamental constant expansions. *Experimental Mathematics*, 10:175, 2001.
- [4] Jonathan M. Borwein, David Borwein, and William F. Galway. Finding and excluding b -ary machin-type BBP formulae. *Available Online*
<http://crd.lbl.gov/~dhbailey/dhbpapers/machin.pdf>, 2002.
- [5] Marc Chamberland. Binary BBP-formulae for logarithms and generalized gaussian-mersenne primes. *Journal of Integer Sequences*, 6, Article 03.3.7, 2003.
- [6] Leonard Lewin. *Polylogarithms and associated functions*. Elsevier North Holland Inc., 1981.
- [7] Jaume Oliver Lafont. Degree 1 BBP-Type Zero Relations. *Private Communication*. Available from the author, January, 2011.